InkChemistry

QUESTION WORKBOOK

Time-of-flight mass spectrometry and relative atomic mass

Atomic structure · Unit C01

AQA 3.1.1.2 · OCR A 2.1.1

14 questions · 46 marks

PRINT EDITION

Every question in this book has a code beside its number.
Type inklearning.co.uk/q/ followed by the code to open that question online.
© 2026 Ink Learning Ltd. All rights reserved.
inklearning.co.uk/chemistry/

SECTION B · Standard, 3 to 5 marks

B1 TFM-Q1

State, in order, the four stages of a time-of-flight mass spectrometer, and explain why the instrument is kept under a high vacuum. [4]

B2 TFM-Q10

Calculate the flight time of a $^{24}Mg^{+}$ ion in a spectrometer that gives every ion 2.00×10^{-16} J of kinetic energy and has a flight tube 1.50 m long. $L = 6.022 \times 10^{23}$ mol^{-1}. [4]

B3 TFM-Q16

Compare electron impact ionisation with electrospray ionisation. [4]

© 2026 Ink Learning Ltd. All rights reserved.

B4 TFM-Q19 Calculate the number of carbon atoms in one molecule of a compound whose mass spectrum has a molecular ion peak of relative height 62.0 and an M+1 peak of relative height 3.41, and explain why the M+1 peak exists. [4]

B5 TFM-Q2 Describe how a sample of krypton is ionised by electron impact, and give an equation for the ionisation. [3]

B6 TFM-Q3 Describe how a sample is ionised by electrospray ionisation, and give the equation for the ionisation of a molecule X. [3]

B7 TFM-Q4 Explain why all the 1+ ions leave the acceleration stage with the same kinetic energy but reach the detector at different times. [3]

© 2026 Ink Learning Ltd. All rights reserved.

B8 TFM-Q5 Explain how the detector of a time-of-flight mass spectrometer produces a signal that measures the abundance of each ion. [3]

B9 TFM-Q9 Calculate the percentage abundance of boron-10 in boron, given that Ar of boron is 10.8 and its only isotopes are boron-10 and boron-11. [3]

B10 TFM-Q11 Calculate the flight time of an $^{81}Br^{+}$ ion in a spectrometer in which a $^{79}Br^{+}$ ion takes 1.54×10^{-5} s, all ions having the same kinetic energy. [3]

B11 TFM-Q14 Suggest the relative molecular mass of a protein whose largest peak in an electrospray spectrum is at m/z = 14306, and suggest why electrospray is used for it rather than electron impact. [3]

© 2026 Ink Learning Ltd. All rights reserved.

B12 TFM-Q15 Suggest why the mass spectrum of sulfur shows a small peak at m/z = 16, although no isotope of sulfur has a mass number of 16. [3]

B13 TFM-Q17 Evaluate the statement that the relative atomic mass of bromine, 79.9, is the mass of a bromine atom. [3]

B14 TFM-Q21 Suggest which halogen is present in a halogenoalkane whose mass spectrum shows two molecular ion peaks of almost equal height, at m/z = 122 and 124, and suggest its molecular formula. [3]

© 2026 Ink Learning Ltd. All rights reserved.

THIS BOOK ONLINE

Scan the code, or type the address, to open the questions in this book online:

inklearning.co.uk/chemistry/practise/questions/time-of-flight-mass-spectrometry/

© 2026 Ink Learning Ltd. All rights reserved.