



ANSWER BOOK

Black-body radiation and spectral classes

Astrophysics · Year 13

AQA 3.9.2.3, 3.9.2.4 · CIE 25.2.1, 25.2.2, 25.2.3 · OCR A 5.5.2(e), 5.5.2(i), 5.5.2(j), 5.5.2(k)

19 questions · 57 marks

NAVY EDITION

SECTION A · Warm-up

- A1** Stefan: $P = \sigma AT^4$, with A the surface area and T the surface temperature in kelvin (1). Wien: $\lambda_{\max} T = 2.9 \times 10^{-3} \text{ m K}$, the constant carrying the product unit metre kelvin (1). [2]
- A2** O B A F G K M, hottest to coolest (1). The Sun, at about 5800 K, is a class G star (1). [2]
- A3** The star radiates as a black body (1), and the light travels to the telescope without absorption on the way, by dust or by atmosphere (1). [2]
- A4** A body that absorbs all the electromagnetic radiation incident on it (1). It is therefore also the best possible emitter, radiating a spectrum that depends only on its temperature (1). [2]
- A5** Class O stars are blue; class M stars are red (1). Class M spectra show molecular absorption by titanium oxide, TiO (1). [2]
- A6** The peak moves to a shorter wavelength (Wien's law) (1), and the curve rises at every wavelength, so the total power radiated (the area under the curve) increases (1). [2]

SECTION B · Standard

- B1** $\lambda_{\max} = 2.9 \times 10^{-3} / 10\,000$ (1) [3]
 $\lambda_{\max} = 2.9 \times 10^{-7} \text{ m}$ (1)
 In the ultraviolet, so the star looks blue-white (1)
- B2** $T = 2.9 \times 10^{-3} / (9.7 \times 10^{-7})$ (1) [3]
 $T = 2990 \text{ K}$ (1)
 Just below 3500 K, so a red class M star (1)
- B3** $A = 4\pi r^2 = 4\pi \times (1.39 \times 10^9)^2$ (1) [4]
 $A = 2.43 \times 10^{19} \text{ m}^2$ (1)
 $P = \sigma AT^4 = 5.67 \times 10^{-8} \times 2.43 \times 10^{19} \times 9000^4$ (1)
 $P = 9.1 \times 10^{27} \text{ W}$ (1)

- B4** Balmer absorption needs hydrogen atoms already excited to the $n = 2$ level (1). [3]
In cool stars almost all atoms sit in the ground state, so the lines are weak; in the hottest stars the hydrogen is ionised, so no bound electrons remain to absorb (1). Class A, 7500 to 11 000 K, is the productive middle (1).
- B5** $P \propto T^4$ at fixed area (1), so the power rises by a factor of $3^4 = 81$ (1). [2]
- B6** $\lambda_{\max}T$ is constant, so the 400 nm star is twice as hot as the 800 nm star (1). With [3]
equal areas $P \propto T^4$ (1), so the ratio is $2^4 = 16$, the hotter star radiating 16 times the power (1).
- B7** $A = 4\pi r^2 = 4\pi \times (7.0 \times 10^6)^2 = 6.2 \times 10^{14} \text{ m}^2$ (1). $T^4 = P/\sigma A$ (1) $= 1.1 \times 10^{24}/(5.67 \times 10^{-8} \times 6.2 \times 10^{14})$ (1), so $T = 1.3 \times 10^4 \text{ K}$ (about 13 000 K) (1). [4]
- B8** Power per unit area $= \sigma T^4 = 5.67 \times 10^{-8} \times 5800^4$ (1) $= 6.4 \times 10^7 \text{ W m}^{-2}$ (1). [2]

SECTION C · Stretch

- C1** $A = P/\sigma T^4 = 3.9 \times 10^{28}/(5.67 \times 10^{-8} \times 3480^4)$ (1) [5]
 $A = 4.7 \times 10^{21} \text{ m}^2$ (1)
 $r = \sqrt{A/4\pi}$ (1)
 $r = 1.9 \times 10^{10} \text{ m}$ (1)
 ≈ 28 solar radii (1)
- C2** X, below 3500 K, peaks towards the infrared and looks red (1); Y, between 7500 [3]
and 11 000 K, peaks towards the ultraviolet and looks blue-white (1). With equal areas, $P \propto T^4$, so Y out-radiates X by a factor of tens to around a hundred (1).
- C3** Measure the wavelength at which the star's black-body spectrum peaks (1); [2]
Wien's law $\lambda_{\max}T = 2.9 \times 10^{-3} \text{ m K}$ then gives the surface temperature directly (1).
- C4** Wien: $T_J = 2.9 \times 10^{-3}/(6.9 \times 10^{-7}) = 4203 \text{ K}$ and $T_K = 12609 \text{ K}$ (1). $A = P/\sigma T^4$ and $r =$ [5]
 $\sqrt{A/4\pi}$ (1). $r_J = 1.4 \times 10^9 \text{ m}$ (1); $r_K = 2.9 \times 10^8 \text{ m}$ (1). Star J is about 5 times larger in radius, despite its smaller power output: its low temperature must be offset by area (1).
- C5** Measure the wavelength at which the star's spectrum peaks; Wien's law $\lambda_{\max}T$ [6]
 $= 2.9 \times 10^{-3} \text{ m K}$ gives the surface temperature (1). Measure the radiant flux at Earth and, knowing the star's distance, use the inverse square law to find the total power radiated (1). Stefan's law $P = \sigma AT^4$ then gives the surface area (1), and $A = 4\pi r^2$ converts area to radius (1). This assumes the star radiates as a black body (1) and that no light is absorbed between the star and the telescope, by dust or atmosphere (1).